

**Jari Hynynen, Pentti Niemistö, Erkki Verkasalo & Anneli Viherä-Aarnio**  
**COST E42 Growing Valuable Broadleaved Tree Species**

**Sept 29, 2006**

**SELECTED FINNISH LITERATURE ON BROADLEAVED TREE SPECIES, UPDATE**

**31. Ecology, genetics and forest tree breeding**

Koski, V. 1991. Experience with genetic improvement of birch in Scandinavia. P. 67-73 in The commercial potential of birch in Scotland, Lorrain-Smith, R., and R. Worrell (eds.). FICGB, London, United Kingdom.

Li, C., Welling, A., Puhakainen, T., Viherä-Aarnio, A., Ernstsen, A., Junntila, O., Heino, P. & Palva, T.E. 2005. Differential responses of silver birch (*Betula pendula*) ecotypes to short day photoperiod and low temperature. *Tree Physiology* 25: 1563-1569.

Oksanen, E., Freiwald, V., Prozherina, N. & Rousi, M. 2005. Photosynthesis of birch (*Betula pendula*) is sensitive to springtime frost and ozone. *Can. J. For. Res.* 35: 703-712.

Oksanen, E., Riikonen, J., Kaakinen, S., Holopainen, T. & Vapaavuori, E. 2005. Structural characteristics and chemical composition of birch (*Betula pendula*) leaves are modified by increasing CO<sub>2</sub> and ozone. *Global Change Biology* 11: 732-748.

Pekkinen, M., Varvio, S., Kulju, K.K.M., Kärkkäinen, H., Smolander, S., Viherä-Aarnio, A., Koski, V. & Sillanpää, J. 2005. Linkage map of birch, *Betula pendula* Roth, based on microsatellites and amplified fragment length polymorphisms. *Genome* 48: 619-625.

Peltonen, P.A., Vapaavuori, E. & Julkunen-Tiitto, R. 2005. Accumulation of phenolic compounds in birch leaves is changed by elevated carbon dioxide and ozone. *Global Change Biology* 11: 1305-1324.

Rousi, M. & Pusenius, J. 2005. Variations in phenology and growth of European white birch (*Betula pendula*) clones. *Tree Physiology* 25(2): 201-210.

Ryynänen, L. & Aronen, T. 2005. Genome fidelity during short- and long-term tissue culture and differentially cryostored meristems of silver birch (*Betula pendula*). *Plant Cell, Tissue and Organ Culture* 83: 21-32.

Ryynänen, L. & Aronen, T. 2005. Vitrification, a complementary cryopreservation method for *Betula pendula* Roth. *Cryobiology* 51: 208-219.

Sillanpää, M., Kontunen-Soppela, S., Luomala, E.-M., Sutinen, S., Kangasjärvi, J., Häggman, H. & Vapaavuori, E. 2005. Expression of senescence-associated genes in the leaves of silver birch (*Betula pendula*). *Tree Physiology* 25: 1161-1172.

Tiimonen, H., Aronen, T., Laakso, T., Saranpää, P., Chiang, V., Ylioja, T., Roininen, H. & Häggman, H. 2005. Does lignin modification affect feeding preference or growth performance of insect herbivores in transgenic silver birch (*Betula pendula*)? *Planta* 222:699-708.

Vakkari, P., Blom, A., Rusanen, M., Raisio, J. and Toivonen, H. 2006. Genetic variability of fragmented stands of pedunculate oak (*Quercus robur*) in Finland. *Genetica* 127:231-241.

Viherä-Aarnio, A., Häkkinen, R. & Junntila, O. 2006. Critical night length for bud set and its variation in two photoperiodic ecotypes of *Betula pendula*. *Tree Physiology* 26: 1013-1018.

Viherä-Aarnio, A., Häkkinen, R., Partanen, J., Luomajoki, A. & Koski, V. 2005. Effects of seed origin and sowing time on timing of height growth cessation of *Betula pendula* seedlings. *Tree Physiology* 25(1): 101-108.

Viherä-Aarnio, A. & Heikkilä, R. 2006. Effect of the latitude of seed origin on moose (*Alces alces*) browsing on silver birch (*Betula pendula*). *Forest Ecology and Management* 229: 325-332.

Viherä-Aarnio, A. & P. Velling, P. 1999. Growth and stem quality of mature birches in a combined species and progeny trial. *Silva Fenn.* 33:225-234.

Ylioja, T. 2000. Relationship between *Phytobia betulae* and its host tree *Betula sp.*. University of Joensuu, Faculty of Forestry, Res. Pap. 112. 26 p. + 5 appendices. Doctoral thesis.

### 32. Silviculture, forest management planning and forest economics

Ahtikoski, A., Päätalo, M-L., Niemistö, P., Karhu, J. & Poutiainen, E. 2004. Effect of alternative thinning intensities on the financial outcome in silver birch (*Betula pendula* Roth) stands: a case study based on long-term experiments and MOTTI stand simulator. *Baltic For.* 10(2): 46-55.

Gustavsen, H.G. & Mielikäinen, K. 1984. Site index curves for natural birch stands in Finland. *Folia For.* 597:1-20. (in Finnish with English summary)

Hynynen, J. & Sanaslahti, A. (eds.). 2002. Management and utilisation of broadleaved tree species in Nordic and Baltic countries – Birch, aspen and alder. Proceedings of the Workshop held in Vantaa, Finland, May 16 to 18, 2001. Finnish For. Res. Inst., Res. Pap. 847: 7-18.

Hynynen, J., Ojansuu, R., Hökkä, H., Siipilehto, J., Salminen, H. & Haapala, P. 2002. Models for predicting stand development in MELA System. Finnish For. Res. Inst., Res. Pap. 835. 116 p.

Kärki, T. 2000. Species, furniture type, and market factors influencing furniture sales in Southern Germany. *For. Prod. J.* 50(4): 85-90.

Kärki, T., Eerikäinen, K., Heinonen, J. & Korhonen, K.T. 1999. [Volume tables of grey alder (*Alnus incana*)]. *Metsätieteen aikakauskirja – Folia Forestalia* 1/1999: 39-49. In Finnish.

Kärki, T., Maltamo, M. & Eerikäinen, K. 2000. Diameter distribution, stem volume and stem quality models for grey alder (*Alnus incana*) in eastern Finland. *New Forests* 20(1): 65-86.

Mäkinen, H., Ojansuu, R. & Niemistö, P. 2003. Predicting external branch characteristics of planted silver birch (*Betula pendula* Roth.) on the basis of routine stand and tree measurements. For. Sci. 49:301-317.

Niemistö, P. 1995a. Influence of initial spacing and row-to-row distance on the crown and branch properties and taper of silver birch (*Betula pendula*). Scand. J. For. Res. 10:235-244.

Niemistö, P. 1995b. Influence of initial spacing and row-to-row distance on the growth and yield of silver birch (*Betula pendula*). Scand. J. For. Res. 10:245-255.

Niemistö, P. 1996. Yield and quality of planted silver birch (*Betula pendula*) in Finland - Preliminary review. Norw. J. Agr. Sc., Suppl. 24: 51-59.

Niemistö, P. & Väärä, T.. (eds.). Silver Birch Today and in the Future. Research Day in Tampere, March 12, 1997. Finnish For. Res. Inst., Res. Pap. 668: 127-140. In Finnish.

Oikarinen, M. 1983. Growth and yield models for silver birch (*Betula pendula*) plantations in southern Finland. Commun. Inst. For. Fenn. 113. 75 p. (in Finnish with English summary)

Raulo, J. 1977. Development of dominant trees in *Betula pendula* Roth and *Betula pubescens* Ehrh. plantations. Seloste: Viljelyjen raudus- ja hieskoivikoiden valtapuiden kehitys. Commun. Inst. For. Fenn. 90(4).

Valkonen, S., Rantala, S. & Sipilä, A. 1995. [Cultivation and management of noble broadleaves and common alder] Finnish For. Res. Inst., Res. Pap. 575. 112 p.

### 33. Wood science and technology

Asikainen, A. 2002. Chemistry and discolouration of birch wood. In: Paavilainen, L. & Marttila, S. (eds.). Finnish Forest Cluster Research Programme WOOD WISDOM (1998-2001). Final report. Report 3/2002.

Hallaksela, A.M. & Niemistö, P. 1998. Stem discoloration of planted silver birch. Scand. J. For. Res. 13:169-176.

Heräjärvi, H. 2001. Internal knottiness with respect to sawing patterns in *Betula pendula* and *B. pubescens*. Baltic For. 8(1): 42-50.

Heräjärvi, H. 2001. Technical properties of mature birch (*Betula pendula* and *B. pubescens*) for saw milling in Finland. Silva Fenn. 35:469-485.

Heräjärvi, H. 2002. Properties of birch (*Betula pendula*, *B. pubescens*) for sawmilling and further processing in Finland. Finnish For. Res. Inst., Res. Pap. 871. 52 p. + App. Doctoral thesis.

Heräjärvi, H. 2004. Static bending properties of Finnish birch wood. Wood Sci. Tech. 37(6): 523-530.

Heräjärvi, H. 2004. Variation of basic density and Brinell hardness within mature *Betula pendula* and *B. pubescens* stems. Wood and Fib. Sci. 36(2): 216-227.

Heräjärvi, H. 2006. Static bending and swelling properties of strand-made beams of *Pinus sylvestris* and *Betula* sp. In: Morlier, P., Morais, J. & Dourado, N. (eds.). Proceedings of the third International Conference of the European Society for Wood Mechanics (ESWM), 5-8 September 2004. UTAD, Vila Real. pp. 3-10.

Heräjärvi, H., Jouhiaho, A., Tammiruusu, V. & Verkasalo, E. 2004. Small-diameter Scots pine and birch timber as raw materials for engineered wood products. Int. J. For. Eng. 15(2): 23-34.

Heräjärvi, H. & Junkkonen, R. 2006. Wood density and growth rate of European and hybrid aspen in southern Finland. Baltic For. 12(1): 2-7.

Heräjärvi, H., Junkkonen, R., Koivunen, H., Metros, J., Piira, T. & Verkasalo, E. 2006. European and hybrid aspen as raw material of sawn timber and mechanical further products. Working Papers of the Finnish Forest Research Institute 31. 102 p. In Finnish.  
<http://www.metla.fi/julkaisut/workingpapers/2006/mwp031.htm>.

Heräjärvi, H. & Verkasalo, E. 2002. Timber grade distribution and relative stumpage value of mature Finnish *Betula pendula* and *B. pubescens* when applying different bucking principles. For. Prod. J. 52(7/8): 40-51.

Heräjärvi, H., Verkasalo, E., Kaurala, H. & Lehtimäki, J. 2000. Properties and utilisation of birch (*Betula pendula*, *B. pubescens*) for saw milling and further processing in Finland. In: Usenius, A. & Kari, P. (eds.). Proceedings of COST Action E10 Wood Properties for Industrial Use, Third Workshop on Measuring of Wood Properties, Grades and Qualities in the Conversion Chains and Global Wood Chain Optimisation, 19<sup>th</sup> – 21<sup>st</sup> June, 2000, Dipoli, Espoo, Finland, pp. 83-102.

Junkkonen, R. & Heräjärvi, H. 2006. Bending strength and fracture of finger-joints of modified and unmodified *Populus* sp. components: a comparison of air-dry and wet wood jointed with six different glues. In: Fioravanti, M. & Maccioni, N. (eds.). Pre-proceedings of the International Conference on Integrated Approach to Wood Structure, Behaviour and Applications. Joint meeting of the European Society for Wood Mechanics and COST Action E35, Florence, Italy, May 15-17. pp. 274-279.

Kärki, T. 1999. Predicting the value of alder logs (*Alnus incana*) based on external quality. Silva Fenn. 33(1): 13-23.

Kärki, T. 2000. Grey alder (*Alnus incana*) as a raw material for mechanical wood processing in Finland. Finnish For. Res. Inst., Res. Pap. 764. 48 p. + App. Doctoral thesis. In Finnish.

Kärki, T. 2001. Variation of wood density and shrinkage in European aspen (*Populus tremula*). Variation der Holzdichte und des Schwindverhaltens in *Populus tremula*. Holz Roh. U. W. 59(1/2): 79-84.

Kärki, T. & Vainikainen, V. 2004. Determining the quality of aspen (*Populus tremula*) logs for mechanical wood processing in Finland. For. Prod. J. 54(7/8): 64-71.

Kaurala, H., Heräjärvi, H. & Verkasalo, E. 2004. [Technical quality of birch from one-species birch stands and mixed stands of spruce and birch in sawmilling] Metsätieteen aikakauskirja – Folia Forestalia 2/2004: 129-143. In Finnish.

Lindblad, J., Tammiruusu, V., Kilpeläinen, H., Lehtimäki, J., Heräjärvi, H. & Verkasalo, E. 2003. [Utilisation of small-diameter birch for the needs of furniture industries. Final report of a research and development project] Finnish For. Res. Inst., Res. Pap. 899. 68 p. + App. In Finnish.

Luostarinen, K. 2006. Effects of environmental and internal factors of trees and timber treatment on colour of dried birch (*Betula pendula*) wood. Dissertationes Forestales 19. 24 p. + App. Doctoral thesis.

Luostarinen, K., Möttönen, V., Asikainen, A. & Luostarinen, J. 2002. Birch (*Betula pendula*) wood discolouration during drying. Effects of environmental factors and wood location in the trunk. Holzforschung 56: 348-354.

Luostarinen K. & Verkasalo E. 2000. Birch as sawn timber and in mechanical further processing in Finland. A literature study. Silva Fenn. Monogr 1. 40 p.

Mononen, K. 2004. Discoloration of silver birch (*Betula pendula*) wood induced by kiln drying and hydrogen peroxide bleaching. University of Joensuu, Department of Chemistry, Publications 69. 41 p. + App. Doctoral thesis.

Möttönen, V. 2006. Variation in drying behaviour and final moisture content of wood during conventional low temperature drying and vacuum drying of *Betula pendula* timber. Drying Technology (in press).

Möttönen, V. 2005. Variation of colour and selected physical and mechanical properties related to artificial drying of sawn silver birch (*Betula pendula* Roth) timber from plantations. Dissertationes Forestales 13. 43 p. + App. Doctoral thesis.

Möttönen, V., Heräjärvi, H., Koivunen, H. & Lindblad, J. 2004. Influence of Felling Season, Drying Method and Within-tree Location on the Brinell Hardness and Equilibrium Moisture Content of Wood from 27-35-year old *Betula pendula*. Scand. J. For. Res. 19: 241-249, 2004.

Möttönen, V. & Luostarinen, K. 2004. Discolouration of sawn birch (*Betula pendula*) timber from plantation forests during drying: Effects of growing site, felling season and storage of logs on discolouration. Baltic For. 10(2): 31-38.

Möttönen, V. & Luostarinen, K. 2006. Variation in density and shrinkage of birch (*Betula pendula*) timber from plantations and naturally regenerated forests. For. Prod. J. (In press).

Nevalainen, S. 2006. Discolouration of birch after sapping. In: Solheim, H. & Hietala, A.M. (eds.). Forest pathology research in the Nordic and Baltic countries 2005. Proceedings from the SNS meeting in Forest Pathology at Skogbrukets Kursinstitutt, Biri, Norway, 28-31 August 2005. Skogforsk, Norsk Institutt for skogforskning. Aktuelt fra skogforskningen 1/06: 32-36.

Piispanen, R. 2004. Factors affecting silver birch (*Betula pendula* Roth) wood structure and chemistry – growth hormones, genes, defoliation, within-stem and seasonal variation. Finnish For. Res. Inst., Res. Pap. 922. 44 p. + App. Doctoral thesis.

Verkasalo, E. 1990. Koivu ja haapa mekaanisen metsäteollisuuden raaka-aineena Yhdysvalloissa. Summary: Birch and Aspen as a Raw Material for Mechanical Forest Industries in the United States. Finnish For. Res. Inst., Res. Pap. 367. 93 p.

Verkasalo, E. 1993. Koivupuutavaran vikaantuminen pitkittyneessä metsävarastoinnissa ja sen vaikutus viilun saantoon, laatuun ja arvoon. Summary: Deterioration of Birch Timber during Prolonged Storage in the Forest and Its Effect on the Yield, Quality and Value of Rotary-Cut Veneer. Folia For. 806. 31 p.

Verkasalo, E. 1997. Quality of European white birch (*Betula pubescens* Ehrh.) for veneer and plywood. Finnish Forest Research Institute, Research Papers 632. 483 p. + App. 59 p. (in Finnish with English summary)

Verkasalo, E., Heräjärvi, H., Arponen, J. & Toppinen, A. 2006. Wood resources and industrial perspectives of plywood industries in the Baltic Sea area. In: Van Acker, J., Irle, M. & Oliver, J.-V. (eds.). Wood Resources and Panel Properties. Conference proceedings, pp. 309-312. Conference co-organized by Cost Action E44 – E49, Valencia, Spain, 12 -13 June 2006. AIDIMA - Furniture, Wood and Packaging Technology Institute.

Verkasalo, E., Sipi, M., Luostarinen, K. & Kärki, T. 2002. Properties of domestic birch, aspen and alder and their utilisation in mechanical wood processing. In: Paavilainen, L. & Marttila, S. (eds.). Finnish Forest Cluster Research Programme WOOD WISDOM (1998-2001). Final Report. Report 3/2002.

Verkasalo, E., Tahvanainen, T., Kilpeläinen, H., Sikanen, L. Heräjärvi, H. & Lindblad, J. 2005. Wood fuels in birch saw mills's product palette in Finland. In: Asplund, D. (ed.). Bioenergy in Wood Industry 2005, 12.-15.9.2005 – Book of Proceedings. FINBIO Publication 32, pp. 243-250.

Viitaniemi, P., Jämsä, S., Paajanen, T., Vuorinen, T., Sundholm, F., Maunu, S. & Paakkari T. 2002. Reaction mechanisms of modified wood. In: Paavilainen, L. & Marttila, S. (eds.). Finnish Forest Cluster Research Programme WOOD WISDOM (1998-2001). Final Report. Report 3/2002.

Zaman, A., Alén, R. & Kotilainen, R. 2000. Thermal behavior of Scots pine (*Pinus sylvestris*) and silver birch (*Betula pendula*) at 200-230°C. Wood Fiber Sci 32:138-143.